

A letter contests the view that a CT scan of *Australopithecus africanus* "is about to wreak havoc on our view of hominid evolution." Other letter writers discuss theropod dinosaur and ancient bird "breathing mechanisms," looking at how "lung ventilation and gas exchange" relate to the dinosaur-origin-of-birds hypothesis.

No Surprises? Dean Falk's Research commentary "Hominid

brain evolution: Looks can be deceiving" (*Science's* Compass, 12 June, p. 1714) contends that "another new specimen...is about to wreak havoc on our view of hominid evolution" and that it changes the big picture. A cranium (Stw 505) found at Sterkfontein, South Africa, in 1989 was no surprise—the site has been producing specimens for more than 50 years. A coffee-table book (1) and "the grapevine" speculated that it had a 600+ cubic-centimeter (cm^3) brain. To test this, brain volume was estimated by using a computed tomography (CT) scanner. The digital measure matched the volume obtained by traditional water displacement, but not the coffee-table book or the grapevine. Conclusion: CT scanners are more accurate than rumors. To have these sources match, or to have found a brain, *would* have been surprising. A mere volume of 515 cm^3 for *Australopithecus africanus* is not. Readers still searching for the surprise, anticipating havoc in the field and changes in their "big picture," may be disappointed by these facts, but not all summer "blockbusters" live up to reviews by friendly critics.

Tim White

Laboratory for Human Evolutionary Studies, Museum of Vertebrate Zoology, and Department of Integrative Biology, University of California, Berkeley, CA 94720, USA, E-mail: timwhite@socrates.berkeley.edu

References

1. D. Johanson and B. Edgar, *From Lucy to Language* (Simon & Schuster, New York, 1996).

Response:

Stw 505 is important because of its unexpected cranial capacity of 515 cm^3 , which suggests that the cranial capacities of other hominid fossil specimens may have been incorrectly estimated. Since writing my commentary, my colleagues and I have extensively researched the endocasts in our collection and have begun to compare them with 3D-CT reconstructions from the actual specimens from which they were taken. We are mighty surprised, and intend

to publish our results as quickly as you can say "*Ardipithecus*." When reading a film review, I always hope that the critic (friendly or otherwise) has actually seen the picture. Worthwhile science, like a credible film review, depends on knowledge of the subject matter. In other words, the proof, as it relates to the significance of Stw 505, is in the pudding. I believe that White and I are both attending a small workshop in August. I'm bringing the pudding. White should bring a spoon.

Dean Falk

Department of Anthropology, State University of New York at Albany, Albany, NY 12222, USA, d.falk@albany.edu

Lung Ventilation and Gas Exchange in Theropod Dinosaurs

John A. Ruben and his colleagues suggest (Reports, 14 Nov. 1997,

p. 1267) that the lung of theropod dinosaurs was most likely similar in form to that of several extant reptiles and was therefore incapable of sustaining high oxygen (O_2) exchange rates characteristic of endothermy. We disagree for two reasons. First, we examined the comparative physiology literature and determined that maximum oxygen exchange rates (VO_2max) of some extant reptiles overlap the oxygen consumption rates measured in some mammals during activity. Specifically, exceptionally active reptiles with multicameral lungs (1) (for example, monitor lizards and sea turtles) have values of VO_2max that overlap or approach the oxygen exchange rates measured in similar size mammals during activity (2). Therefore, the septate lung in those reptiles must be capable of sustaining rates of gas flux characteristic of endotherms. However, mammals and birds "typically" have a greater VO_2max . Therefore, we addressed the question of what modifications in the oxygen transport system of an extant reptile would be necessary to support higher rates of oxygen consumption.

We used morphological and physiologi-

cal measurements of extant reptiles and well-established respiratory equations to model the gas exchange potential of the reptilian oxygen delivery system and to examine the role of lung structure in constraining gas exchange. Each step in the oxygen cascade is described by a set of respiratory equations and, consequently, it is possible to describe mathematically the flux of oxygen through the entire cascade and to evaluate the impact of modifications in any of its components (3). We used this approach to predict the effects of modifying several parameters in the oxygen cascade on VO_2max in a 1-kilogram lizard, *Varanus exanthematicus* (4). Our analysis included four modifications: (i) a small increase in the maximum cardiac output; (ii) an increased oxygen carrying capacity of the blood from reptilian to mammalian values; (iii) an increase in maximum cardiac output combined with the changes in blood oxygen-carrying capacity; and (iv) an increased respiratory gas exchange area in the dorsal region of the lung through elaboration of the intercameral septa with a membranous region in the ventral portion of the lung. Without modification of the lung structure, our analysis predicts that changes in blood oxygen capacity and cardiac output support a VO_2max that is 50% of the value for a "typical" 1-kilogram (kg) mammal (5). However, if we combine these changes with conservative modifications in lung morphology, we predict a VO_2max that is nearly 70% of the typical mammalian value. Our analysis indicates that modifications in several of the steps of the oxygen cascade have a cumulative effect on VO_2max (6). The resulting high oxygen flux rate mandates an increase in lung ventilation that is 233% above the maximum level measured in extant lizards.

Lizards have a mechanical constraint on simultaneous vigorous locomotion and costal ventilation that arises from the design of the axial musculoskeletal system, and this mechanical constraint was probably the primitive condition for all tetrapods (7). Consequently, the fundamental change required to support sustainable high oxygen exchange rates was the development of new mechanisms to increase ventilation (7). This constraint has been circumvented to varying degrees in some extant lizards, for example, the use of the gular pump to assist costal ventilation during activity (8) and in the lineages that gave rise to endotherms by the evolution of ventilatory mechanics that are not limited by locomotor requirements (7).

Inadequate preservation of the soft-tissue components of the oxygen transport

system precludes accurate assessment of the aerobic potential of theropod dinosaurs. However, on the basis of metabolic patterns in extant reptiles and our theoretical analysis, we find that the notion that nonavian septate lungs constrain high oxygen flux rates is not supported. Our analysis suggests that modifications in lung structure were not a prerequisite for supporting higher oxygen consumption rates. In the mammalian and archosaur lineages that evolved endothermy, higher oxygen consumption rates could have been supported through changes in ventilatory mechanics and increases in blood oxygen content and cardiac output.

James W. Hicks

Colleen G. Farmer

Department of Ecology and Evolutionary Biology,
University of California, Irvine, CA 92697, USA

References

1. S. F. Perry, *Adv. Anat. Embryol. Cell Biol.* **79**, 1 (1983).
2. A. W. Crompton, C. R. Taylor, J. A. Jagger, *Nature* **272**, 333 (1978); C. R. Taylor, N. C. Heglund, G. M. O. Maloiy, *J. Exp. Biol.* **97**, 1 (1982); G. S. Mitchell, T. T. Gleeson, A. F. Bennett, *Respir. Physiol.* **43**, 365 (1981); D. Jackson and H. D. Prange, *J. Comp. Physiol.* **134**, 315 (1979); F. V. Paladino, M. P. O'Connor, J. R. Spotila, *Nature* **344**, 858 (1990).
3. E. R. Weibel, *The Pathway for Oxygen* (Harvard Univ. Press, Cambridge, MA, 1984); P. D. Wagner, *Respir. Physiol.* **106**, 329 (1996).

4. J. W. Hicks and C. W. Farmer, in preparation.
5. C. R. Taylor *et al.*, *Respir. Physiol.* **44**, 25 (1981).
6. Table 1 at <http://compphys.bio.uci.edu/hicks/lung.htm>
7. D. R. Carrier, *Paleobiology* **13**, 326 (1987).
8. E. L. Brainerd and T. Owerkowicz, *Am. Zool.* **36**, 88A (1996).

Ruben *et al.* are to be commended for bringing forth yet another intriguing argument concerning dinosaur physiology. While there may be evidence that theropod dinosaurs maintained a separate thoracic and abdominal cavity by means of a septum, stronger evidence is needed to support the idea that this membrane served as a diaphragmatic hepatic piston. The authors compare by analogy the partitioning of the body cavity of an alligator with *Sinosauropteryx* (their figure 5). It would be beneficial to show in a similar illustration how this condition differs in other reptiles, as well as in birds. Teiid lizards (such as *Tupinambis*), for example, independently developed a non-muscular posthepatic septum dividing the lungs and liver from the abdominal viscera (1), yet unlike in crocodilians, this structure does not participate in ventilation. Varanid lizards (such as *Varanus*) similarly separate their lungs (which occupy most of the thorax)

from the peritoneal cavity by means of a non-ventilatory, postpulmonary septum (1). Birds also developed a complex, non-muscular, septal partitioning of their body cavity by means of two membranes: the horizontal and the oblique septa (the latter separates all but the abdominal air sacs from the viscera, spanning the vertebrae to the sternal margin) (2). These septa, also, are nonventilatory in function. How can we rule out that we are not seeing something similar in *Sinosauropteryx*?

Ruben *et al.* also note the lack of an "expansive sternum," "hinged ribs," and evidence for "inhalatory filling of the abdominal air sacs" as indicators that theropods lacked modern avian lungs. The primary reason for the elongation of the sternum, however (along with the development of the sternal carina), appears to have been to increase the surface area for the origin of the flight muscles. The hinged ribcage can also be considered an adaptation to counter the forces incurred by the thorax during flight. What is the empirical evidence to argue that these features must have been in place before a modern avian lung could have developed? Additionally, the lack of air sac diverticula into bone may not have precluded the possibility of

NEW!


~~quick as greased lightning~~

~~pronto~~ ~~in a jiffy~~ ~~lickity-split~~

~~immediate gratification~~

There's only one way to say it...

Now you can order FREE product literature instantly on the Internet with SCIENCE's Literature Library. Whether you're looking for the latest catalogs, product brochures, or technical application sheets, you'll find them all here. Best of all, you'll be able to request information directly from the world's leading life science suppliers with the Literature Library's e-mail order form.



SCIENCE's Literature Library

www.sciencemag.org

Click on Electronic Marketplace, then click on Literature Library.